

Bimectin Injection

Ivermectin 1% w/v

INDICATIONS

Contains 1% ivermectin w/v
Licenced for Cattle, Sheep and Pigs.

Cattle

Licensed for wide range of internal and external Parasites.

Sheep

Licensed for a wide range of Gastro-Intestinal worms, Lungworms and Nasal Bots.

Pigs

Licensed for a wide range of external and Internal parasites.

BENEFITS

- ✓ Antiparastic injectable solution
- ✓ Delivers effective control against a wide range of external and internal parasites
- ✓ Penetrates quickly to reach and kill parasites
- ✓ Licenced for cattle, sheep and pigs
- ✓ Cost effective parasitic control in animals
- ✓ 1 x 500ml pack can treat; 333 x 50-75kg ewes, or 62 x 400kg cattle, or 125 x 100- 133kg pigs



PACKAGING

List No	Unit Package	Case Size
1BIM075	250ml	6
1BIM076	500ml	6



Bimectin Injection

Ivermectin 1% w/v



PRESENTATION

A clear, colourless slightly viscous, non-aqueous sterile solution containing 1% w/v Ivermectin.

TARGET SPECIES

Cattle, Sheep and Pigs.

INDICATIONS FOR USE, SPECIFYING THE TARGET SPECIES

For the effective treatment and control of the following harmful parasites of cattle, sheep and pigs:

Cattle

Gastrointestinal roundworms (adult and fourth-stage larvae):

Ostertagia spp. (including inhibited *O. ostertagi*)

Haemonchus placei

Trichostrongylus axei

T. colubriformis

Cooperia spp.

Bunostomum phlebotomum

Oesophagostomum radiatum

Strongyloides papillosus (adult)

Nematodirus helvetianus (adult)

N. spathiger (adult)

Trichuris spp (adult).

Lungworms (adult and fourth-stage larvae):

Dictyocaulus viviparus

Eye worms(adult):

Thelazia spp.

Warbles:

Hypoderma bovis

H. lineatum

Mange mites:

Psoroptes bovis

Sarcoptes scabiei var. *bovis*

Suckling lice:

Linognathus vituli

Haematopinus eurysternus

Solenopotes capillatus

May also be used as an aid in the control of the mange mite *Chorioptes bovis* and biting lice *Damalinea bovis*, but complete elimination may not occur.

Persistent Activity

Treatment at the recommended dose rate can control re-infection with *Haemonchus placei* and *Cooperia* spp. acquired up to 14 days after treatment, *Ostertagia ostertagi* and *Oesophagostomum radiatum* acquired up to 21 days after treatment and *Dictyocaulus viviparus* up to 28 days after treatment.

To obtain optimal benefit from the persistent activity of the product for grazing animals, it is recommended that calves which are set-stocked in the first grazing season should be treated 3, 8 and 13 weeks after the day of turn-out. This can protect the animals from parasitic gastroenteritis and lungworm disease throughout the grazing season, provided they are set-stocked, all the calves included in the programme and that no untreated cattle are added to the pasture. Treated animals should always be monitored according to good husbandry practices.

Sheep

Gastrointestinal roundworms (adult and fourth-stage larvae):

Ostertagia circumcincta including inhibited larvae

O. trifurcata

Haemonchus contortus including inhibited larvae

Trichostrongylus axei (adults)

T. colubriformis and *T. vitrinus* (adults)

Cooperia curticei

Oesophagostomum columbianum

O. venulosum (adults)

Nematodirus filicollis

Chabertia ovina

Trichuris ovis (adults).

Lungworms:

Dictyocaulus filaria (adult and fourth-stage larvae)

Protostrongylus rufescens (adults)

Nasal Bots (all larval stages)

Oestrus ovis

Pigs

Gastrointestinal roundworms (adult and fourth-stage larvae):

Ascaris suum

Hyostrongylus rubidus

Oesophagostomum spp.

Strongyloides ransomi (adult and somatic larval stages)

Lungworms:

Metastrongylus spp. (adults)

Lice:

Haematopinus suis

Mange mites:

Sarcoptes scabiei var. *suis*

CONTRAINDICATIONS

Do not use in animals with known hypersensitivity to the active ingredient.

Do not use by intramuscular or intravenous administration.

SPECIAL WARNINGS FOR TARGET SPECIES

None

SPECIAL PRECAUTIONS FOR USE

(i) Special Precautions for use in animals

The product has been formulated specifically for use in cattle, sheep and pigs. It should not be used in other species as severe adverse reactions, including fatalities in dogs, may occur.

Frequent and repeated use may lead to the development of resistance. It is important that the correct dose is given in order to minimise the risk of resistance. To avoid under-dosing, animals should be grouped according to their bodyweight and dosed according to the dose of the heaviest animal in the group.

(ii) Special Precautions to be taken by the Person Administering the Product to Animals

Take care to avoid self-administration: the product may cause local irritation and/or pain at the site of injection.

Direct contact of the product with the skin should be kept to a minimum.

Do not smoke or eat while handling the product. Wash hands after use.

(iii) Other precautions

When using the 250 ml and 500 ml pack sizes, use only automatic syringe equipment. To refill the syringe, use of a draw off needle is recommended to avoid excessive broaching of the stopper.

ADVERSE REACTIONS (FREQUENCY AND SERIOUSNESS)

Cattle

Transient discomfort has occasionally been observed in cattle following subcutaneous administration. A low incidence of soft tissue swelling at the injection site has been observed.

Sheep

Discomfort, sometimes intense but usually transient, has been observed in some sheep immediately following subcutaneous administration.

Pigs

Mild and transient discomfort has occasionally been observed in pigs following subcutaneous injection.

All these reactions disappeared without treatment.

USE DURING PREGNANCY AND LACTATION OR LAY

The product can be administered to beef cows, sheep and pigs at any stage of pregnancy

Lactation

Do not use in dairy cows or sheep producing milk for human consumption.

Do not use in non-lactating dairy cows or sheep within 60 days of calving/lambing. The product can be used in sows during lactation.

Fertility

Fertility is not affected by administration of the product.

INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION

The product can be used concurrently without adverse effects with foot and mouth disease vaccine or clostridial vaccine, given at separate injection sites.

AMOUNTS TO BE ADMINISTERED AND ADMINISTRATION ROUTE

The product should be given only by subcutaneous injection at the recommended dosage level of 200 mcg ivermectin per kg bodyweight under the loose skin in front of, or behind, the shoulder in cattle and over the neck in sheep. At the recommended dosage level of 300 mcg ivermectin per kg of bodyweight, the product should be given only subcutaneously in the neck of pigs.

Each ml contains 10 mg of ivermectin sufficient to treat 50 kg of bodyweight of cattle and sheep and 33 kg of bodyweight of pigs.

The injection may be given with any standard automatic or single-dose or hypodermic syringe.

Use of a sterile 17 gauge x 1/2 inch needle is suggested.

Injection of wet or dirty animals is not recommended.

If using a single-dose or hypodermic syringe, use a separate sterile needle to withdraw the product from the container.

Massage the injection site after administration of the product.

In young pigs, especially those below 16 kg for which less than 0.5 ml of the product is indicated, dosing accuracy is important. The use of a syringe that can accurately deliver as little as 0.1 ml is recommended.

In young lambs weighing less than 20.0 kg give 0.1 ml per 5 kg. In these lambs the use of a syringe with can deliver as little as 0.1 ml is recommended.

OVERDOSE (SYMPTOMS, EMERGENCY PROCEDURES, ANTIDOTES), IF NECESSARY

Cattle

Single doses of 4.0 mg ivermectin per kg (20 x the use level) given subcutaneously resulted in ataxia and depression.

Sheep

At dose levels up to 4 mg ivermectin per kg (20 x the use level) given subcutaneously resulted in ataxia and depression.

No signs of systemic toxicity were observed in sheep treated with the product at up to 3 times the recommended dose rate, soft tissue swellings at the injection site were observed.

Pigs

A dose of 30 mg ivermectin per kg (100 x the recommended dose of 0.3 mg per kg) injected subcutaneously to pigs caused lethargy, ataxia, bilateral mydriasis, intermittent tremors, laboured breathing and lateral recumbency.

In the case of overdose, symptomatic treatment should be given.

WITHDRAWAL PERIOD(S)

Cattle Meat and Offal – 49 days. Do not use in lactating cows producing milk for human consumption. Do not use in non-lactating dairy cows including pregnant dairy heifers within 60 days of calving.

Sheep must not be treated within 42 days of slaughter for human consumption. Do not use in lactating sheep producing milk for human consumption. Do not use in sheep within 60 days of lambing where milk is to be used for human consumption.

Pigs must not be treated within 28 days of slaughter for human consumption.

INCOMPATIBILITIES

Do not mix with other medicinal products.

SHELF LIFE

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years

Shelf-life after first opening the immediate packaging: 28 days.

SPECIAL PRECAUTIONS FOR STORAGE

None

SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED VETERINARY MEDICINAL PRODUCT OR WASTE MATERIAL, IF ANY

Any unused veterinary product or waste material derived from the product should be disposed of in accordance with local requirements. The product should not enter water courses as this may be dangerous to fish and other aquatic organisms.

MARKETING AUTHORISATION HOLDER:

Bimeda Chemicals Ltd., Broomhill Road, Tallaght, Dublin 24, Ireland.

MARKETING AUTHORISATION NUMBER:

VM 12597/4029

LEGAL CATEGORY

POM-VPS

PACKAGE QUANTITIES

250ml, 500ml

A full product SPC is available on request from Bimeda or alternatively can be found on the VMD website