

Novaquin

15mg/ml meloxicam
oral suspension
for horses

INDICATIONS

Alleviation of inflammation and relief of pain in both acute and chronic musculo-skeletal disorders in horses.

BENEFITS

- Acute and chronic pain relief
- Can be administered with food or directly into the mouth
- Once daily dosing



PACKAGING

List No	Unit Package
1NOV001	125ml
1NOV002	336ml



Novaquin

15mg/ml meloxicam oral suspension for horses

Presentation

Oral suspension. Yellowish-green viscous oral suspension.

Target Species

Horses

Indications:

Alleviation of inflammation and relief of pain in both acute and chronic musculo-skeletal disorders in horses.

Amounts to be administered and administration route:

To be administered either mixed with food or directly into the mouth at a dosage of 0.6 mg/kg body weight, once daily, up to 14 days. In case the product is mixed with food, it should be added to a small quantity of food, prior to feeding.

The suspension should be given using the measuring syringe provided in the package. The syringe fits onto the bottle and has a kg-body weight scale.

Shake vigorously at least 20 times before use.

After administration of the veterinary medicinal product, close the bottle by replacing the cap, wash the measuring syringe with warm water and let it dry. Avoid introduction of contamination during use.

Contra-indications & Warnings

Do not use in pregnant or lactating mares. Do not use in horses suffering from gastrointestinal disorders such as irritation and haemorrhage, impaired hepatic, cardiac or renal function and haemorrhagic disorders. Do not use in case of hypersensitivity to the active substance or to any of the excipients. Do not use in horses less than 6 weeks of age.

Special precautions for use in animals

Avoid use in any dehydrated, hypovolaemic or hypotensive animals as there is a potential risk of renal toxicity.

Special precautions to be taken by the person administering the veterinary medicinal product to animals.

People with known hypersensitivity to Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) should avoid contact with the veterinary medicinal product. In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Use during Pregnancy or Lactation

Laboratory studies in cattle have not provided any evidence for teratogenic, foetotoxic, or maternotoxic effects. However, no data has been generated for horses. Therefore the use in this species is not recommended during pregnancy and lactation.

Adverse Reactions

(Frequency and Seriousness)
Isolated cases of adverse reactions typically associated with NSAIDs were observed in clinical trials (slight urticaria, diarrhoea). Symptoms were reversible. In very rare cases loss of appetite, lethargy, abdominal pain and colitis have been reported.

In very rare cases anaphylactoid reactions, which may be serious (including fatal), may occur and should be treated symptomatically. If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

The frequency of adverse reactions is defined using the following convention:
- very common (more than 1 in 10 animals displaying adverse reactions during the course of one treatment) - common (more than 1 but less than 10 animals in 100 animals) - uncommon (more than 1 but less than 10 animals in 1,000 animals) - rare (more than 1 but less than 10 animals in 10,000 animals) - very rare (less than 1 animal in 10,000 animals, including isolated reports).

Withdrawal Periods

Meat and offal: 3 days.

Not authorised for use in horses producing milk for human consumption.

Legal Category

POM-V

Marketing Authorisation Holder

Le Vet Beheer B.V. Wilgenweg 7 3421 TV Oudewater The Netherlands

Marketing Authorisation Number

EU/2/15/186/001-002

TAKE TIME



OBSERVE LABEL
DIRECTIONS

Distributed in the UK by

Bimeda, Unit 2/Bryn Cefni Industrial Park, Llangefni LL77 7XA, United Kingdom
Call Bimeda on 0044 (1)248 725 400
www.bimeda.co.uk