

Zero

(Cyromazine 2% w/w soluble granule (SG) formulation)
Insecticide for Professional Use

INDICATIONS

ZERO is a larvicide used to control flies and fly larvae in intensive livestock units, manure heaps and landfill sites. ZERO contains a potent growth disruptor specific to all fly species.

BENEFITS

- ✓ Prevents larvae from developing in to flies
- ✓ Reduces overall fly challenge
- ✓ Infectious disease vectors are reduced in number
- ✓ Reduces stress caused by nuisance flies



PACKAGING

List No	Unit Package
1ZER007	25kg



Zero

(Cyromazine 2% w/w soluble granule (SG) formulation)
Insecticide for Professional Use



PRESENTATION

Cyromazine 2.0% w/w (20g/kg) Soluble Granule (SG) formulation.

INDICATIONS FOR USE

ZERO is a larvicide used to control flies and fly larvae in intensive livestock units, manure heaps and landfill sites.

It can be used in cattle, horse, pig, poultry, goat sheds, rabbit farms and zoological environments.

DIRECTIONS FOR USE

For the Control of Fly Larvae:

Apply ZERO at Intervals of 6 weeks

Apply ZERO at a dose of 250g/10m², independent of the method of application.

Apply ZERO as either dry granules by dispersal if the manure is liquid or very moist, or dilute 250g in 1-4L of water for spraying or 10L of water for pouring using a watering can. If the manure is dry apply by pouring only.

The simultaneous use of an insecticide specific to adult flies, particularly coinciding with the first application of ZERO, can significantly enhance fly control.

In Rearing Facilities for Cattle and Horses

Apply ZERO in a 50cm band along the walls, under and next to machinery installations by pouring or spraying. When treating liquid manure the granules can be dispersed directly. The first application is best carried out 1-3 days after dung removal. In cow sheds with slatted floors apply ZERO uniformly to the entire surface.

In Pig Sties with completely slatted floors apply ZERO to the entire surface. In the case of partially slatted floors apply to the slatted sections and the borders of the rest areas. Apply either by dispersal of dry granules, or pouring, or spraying. If possible treat before housing the pigs; shortly after manure or slurry removal is best.

In Poultry Houses with a slurry reservoir and in the presence of moist or liquid manure, disperse ZERO granules directly. In dry manure situations use the pouring or spraying method.

Apply to the entire surface. Commence treatment approximately 1 week after removal of excrement but before a depth of 10cm has accumulated.

ZERO can also be used in sheep and goat sheds, rabbit farms and zoological gardens in the same manner.

MANUFACTURER

Hokochemie GmbH,
Pannerhofstrasse 7,
CH-6353,
Weggis.
Tel: +41 41390 3188
Fax: +41 41390 3189

MARKETING COMPANY:

Hockley International Ltd,
Hockley House,
3 Longstone Road,
Ashbrook Office Park,
Manchester,
M22 5LB.
United Kingdom
Tel: +44 (0) 161 209 7400
Fax: +44 (0) 161 209 7401
www.hockley.co.uk

DISTRIBUTOR

Bimeda,
Bryn Cefni Industrial Park,
Llangefni,
Anglesey,
LL77 7XA.
Tel: +44 (0) 1248 725 400.
www.bimeda.co.uk

WARNINGS

H412 Harmful to aquatic life with long lasting effects.

P273 Avoid release to the environment.

P280 Wear protective clothing and gloves.

P501 Dispose of contents/container in accordance with national regulations.

- The COSHH (Control of Substances Hazardous to Health) Regulations 2002 (as amended) apply to the use of this product at work.
- Avoid excessive contamination of coveralls.
- Do not contaminate foodstuffs, eating utensils or food contact surfaces.
- This material and its container must be disposed of in a safe way.
- Do not apply directly to livestock and poultry.

- Exclude all persons and animals during treatment.
- Cover all water storage tanks before applications.
- **Hazardous to bees**
- Protect milk machinery and milk containers from contamination.
- Wash hands and exposed skin before meals and after use.
- Keep in a safe place.
- Collect eggs and remove feed, exposed water and milk before application.
- Not to be used on manure intended for the production of mushroom compost.
- Applications should be made to moist areas where flies are breeding.
- In deep pit poultry houses, with vertical multi-tiered cages with automatic scrapers, more effective distribution of ZERO can be achieved by applying a series of reduced doses over a period of 3 weeks. Apply 25g ZERO dissolved in 1 or more litres of water, 3 times weekly on alternate weekdays for a 21 day period, ie a total of 9 applications in a treatment cycle.
- To avoid risks to man and the environment, comply with the instructions for use.

STATUTORY CONDITIONS OF APPROVAL RELATING TO USE.

**FOR USE ONLY AS AN INSECTICIDE.
FOR USE ONLY BY PROFESSIONAL OPERATORS.**

Application rate: See "DIRECTIONS FOR USE" for details on dose rates.

Users must comply in full with all their legal responsibilities under the Control of Pesticides Regulations 1986 (as amended), including complying with all Conditions of Approval.

Read all precautions before use.

HSE No.10127.

USE ONLY IN ACCORDANCE WITH THE DIRECTIONS ON THE LABEL.

TO AVOID RISKS TO MAN AND THE ENVIRONMENT, COMPLY WITH THE INSTRUCTIONS FOR USE

ZERO is a registered community trademark of HokoChemie GmbH.

Use Medicines Responsibly. Noah.co.uk/responsible

Bimeda.co.uk

TAKE TIME



OBSERVE LABEL DIRECTIONS