

ECTOFLY

Cypermethrin - 1.25% w/v Pour-On Solution For Sheep

DATA
SHEET



INDICATIONS:

Correct Usage:

- For the treatment and control of headflies use T bar nozzle (red)
- For the treatment of ticks and biting lice in sheep use straight nozzle (green)
- For the prevention of blowfly strike use the fanspray nozzle (blue)
- For the treatment of blowfly strike use T bar nozzle (red)

BENEFITS

- 6 – 8 week blowfly strike prevention;
- Treats and prevents blowfly strike
- Treats lice
- Treats ticks
- Treats and prevents headflies
- Proven formulation
- Short 8 day meat withhold period
- Can be used all year round
- Contains 48 hour blue dye

LIST No	UNIT PACKAGE	CASE SIZE
1ECT021	2.5L	5
1ECT004	5L	4
1ECT032	20L*	1

*4 x 5L + FREE applicator

See reverse for Administration & Dosage



Ectofly

Cypermethrin

1.25% w/v Pour-On Solution For Sheep



PRESENTATION

A clear blue non-aqueous solution containing 1.25%w/v High Cis (80:20) cypermethrin.

TARGET SPECIES

Sheep.

USES

For the treatment and control of headflies, and treatment of ticks and biting lice in sheep. For the prevention and treatment of blowfly strike in sheep.

AMOUNTS TO BE ADMINISTERED AND ADMINISTRATION ROUTE

On completion of each treatment cycle of the product and before the dosing gun is stored, immediately expel any remaining product in the dosing gun. Administer by means of the recommended applicator gun as follows:

Blowflies:


TREATMENT OF BLOWFLY STRIKE

Use Red T-Bar Nozzle

Apply directly to all the affected parts at the rate of 2.5ml per 100 -150cm² (roughly the size of a hand). The T-bar nozzle should be used. Most strikes will require 5-10ml. Ensure that all affected parts are treated. A single application is sufficient to ensure that larvae are expelled and killed within a few hours.

PREVENTION OF BLOWFLY STRIKE

Use Blue Fan-Spray Nozzle



BODYWEIGHT	DOSE	DOSES PER PACK
Up to 25 kg*	20 ml	50
25-40 kg	30 ml	33
Over 40 kg	40 ml	25

*Do not administer to animals of less than 12.5kg bodyweight.

Apply as a fan-spray to the surface of the fleece on the back and hindquarters of the sheep using the Pour-On Gun fitted with the Fan-Spray Nozzle.

Half the dose should be applied to the shoulders, back and flanks, and half to the rump. The distance between the nozzle and the fleece should be approximately 20cm.

Each dose will require 2 or 3 sweeps to apply. THE PRODUCT WILL PREVENT BLOWFLY STRIKE ONLY ON AREAS COVERED BY SPRAY. THEREFORE IT IS IMPERATIVE TO USE CORRECT GUN AND BLUE FAN-SPRAY NOZZLE AND ENSURE A WIDE AREA OF

COVERAGE BY KEEPING THE NOZZLE APPROX 20CM FROM THE FLEECE WHEN APPLYING.


One application will give 6-8 weeks protection. Re-treat as required.

Lice.

Use Green Straight Nozzle

5ml per 20kg bodyweight up to a maximum of 20ml:

BODYWEIGHT	DOSE	DOSES PER PACK
Up to 20 kg	5 ml	200
21-40 kg	10 ml	100
41-60 kg	15 ml	66
Over 60 kg	20 ml	50




Using the Straight Nozzle on the applicator gun, apply as a pin-stream from the shoulders to the rump along the middle of the backline. Sheep may be treated off-shears or at any time during the year.

Ticks.

Use Green Straight Nozzle

Adult sheep and lambs greater than 10kg: 10ml per 20kg bodyweight, up to a maximum of 40ml. Lambs under 10kg: 5ml followed 3 weeks later by a 10ml application.

BODYWEIGHT	DOSE	DOSES PER PACK
Up to 10 kg	5 ml	200
11-20 kg	10 ml	100
21-40 kg	20 ml	50
41-60 kg	30 ml	33
Over 60 kg	40 ml	25



Using the Straight Nozzle on the Pour-On Gun, apply as a pin-stream from the crown of the head to the top of the rump.


Headflies.

Use Red T-Bar Nozzle

5ml per animal irrespective of size: Apply to the top of the head between the ears using the T-Bar Nozzle.

Take care not to apply product in the sheep's eyes. One application before the start of the headfly season will give up to 4 weeks protection.

Re-treat as required.



This product contains a blue dye to aid in the identification of treated animals. The colour is temporary and will fade from the fleece within 48 hours of application.

CONTRA-INDICATIONS & WARNINGS

Not to be used in lambs less than one week old. For the prevention of blowfly strike. Do not administer to animals of less than 12.5kg bodyweight. Avoid treating very young lambs if the weather is unseasonably hot.

SPECIAL WARNINGS FOR EACH TARGET SPECIES

Ectofly 1.25% w/v Pour-On minimises mothering-up problems after treatment.

However, care must be taken not to apply the product right to the lamb's tail, as a ewe recognises a young lamb partly by the smell of the tail area.

Blowflies are attracted by dirty, damp wool or open

wounds. The more common site for fly strike is the rump due to soiling caused by diarrhoea and urine staining. The efficacy of Ectofly 1.25% w/v Pour-On is reduced in the presence of soiled or dirty wool.

Therefore, sheep must be crutched (dagged) regularly and appropriate worm control measures employed.

Take care not to apply product in the sheep's eyes. This precaution should be observed particularly when treating breeds with little wool on their heads.

Full operating and maintenance instructions, including details on the use of nozzles are supplied with each applicator gun. Please read carefully before use. Maintain applicator gun carefully to ensure accurate dosage.

For external use only.

SPECIAL PRECAUTIONS FOR USE

Make sure when attaching the recommended applicator gun onto the container that the coiled retainer is secured onto the cap and the applicator.

Wear eye protection, protective clothing, rubber gloves and boots when applying the product. Wear a dust mask when applying as a fan-spray for the prevention of blowfly strike.

Wash splashes from skin and eyes immediately with plenty of clean water. If irritation persists seek medical advice immediately.

Remove contaminated clothing immediately and wash exposed skin with plenty of clean water.

Do not eat, drink or smoke whilst using the product. Wash hands and exposed skin before eating, drinking or smoking and after work. Use in a well ventilated area and avoid inhaling the vapour.

ADVERSE REACTIONS (FREQUENCY AND SERIOUSNESS)

None known.

USE DURING PREGNANCY AND LACTATION

No restrictions. Pregnant ewes must be handled with care.

WITHDRAWAL PERIOD(S)

Meat and offal: 8 days.

Milk: 5 days.

INCOMPATIBILITIES

Do not mix with other medicinal products.

LEGAL CATEGORY

POM-VPS

MARKETING AUTHORISATION NUMBER

VM 50146/4017

MARKETING AUTHORISATION HOLDER

Bimeda Animal Health Limited
2, 3 & 4 Airton Close, Airton Road
Tallaght, Dublin 24, Ireland
Tel: +353 (0) 1 466 7900

DISTRIBUTOR

Bimeda UK
Unit 2, Bryn Cefni Industrial Park
Llangefni, Anglesey,
Wales, LL777XA
Tel: +44 (0) 1248 725 400

Use Medicines Responsibly.
Noah.co.uk/responsible

TAKE TIME



OBSERVE LABEL DIRECTIONS

www.bimeda.co.uk